



Organza 0-0-30+TE

Complete high potassium fertilizer containing organic acids and amino acids

Formulation: Gravel

Saturation level: 212 grams per liter



HANNANN NAHADE ASIA

SOIL REGENERATION ,
SUSTAINABLE CULTIVATION

Introduction

NPK0-0-30+TE of HANNANN Company contains high amounts of potassium and organic compounds such as amino acid and fulvic acid. Also, this fertilizer is rich in micronutrient elements, which increases the absorption and then improves the quantitative and qualitative performance of the products. The formulation of this fertilizer is in the form of granules and it is produced through special mechanisms in such a way that it is completely soluble in water. Also, with high amounts of potassium, this fertilizer has an effective role in improving the quantitative and qualitative performance of agricultural and garden crops.

Benefits and Particulars

- Extremely fast solubility in water
- Can be used as a topical positioning spray and with irrigation water
- Increasing absorption of nutrients
- Helping to improve soil health by reducing the use of chemical fertilizers
- Increasing the activity of soil microorganisms
- Increasing plant resistance against environmental stresses and pathogens
- Increasing the quantitative and qualitative performance of products
- Contains a high Percentage Of Potassium to ripen the fruit and increase the quality of the product
- Eco Friendly

| Element | Percent | Element | Percent |
|----------------------------|---------|-------------------|---------|
| Potassium soluble in water | 30 | Free Amino Acid | 0.2 |
| Soluble Fe | 0.01 | Soluble Zn | 0.002 |
| Soluble Cu | 0.0005 | Fulvic Acid | 3 |
| Soluble Magnesium | 9 | Organic materials | 6 |

| Product | Method of Use | Amount of Use |
|--------------|---------------------|--|
| Garden | Topical Positioning | 0.3-0.5 Kg for every tree (50 gr Per Year Of Tree Age) |
| | Irrigation water | 15-20 Kg in hectare (2 times) |
| | Spraying | 2-3 Kg in each 1000 Liter of water |
| Agricultural | Under cultivation | 50-100 Kg in hectare |
| | Irrigation water | 10-15 Kg in hectare |
| | Spraying | 1-2 Kg in each 1000 Liter of water |

Notes:

- Avoid mixing with very acidic and alkaline compounds.
- Spraying should be done early in the morning or at sunset.
- In case of mixing with other compounds and poisons, it should be tested first in a small amount and at a limited level.